

Unit**06****ALGEBRAIC MANIPULATION****Highest Common Factor (H.C.F.)**

If two or more algebraic expressions are given then their common factor of highest power is called the H.C.F of the expressions.

Least Common Multiple (L.C.M)

If an algebraic expression $p(x)$ is exactly divisible by two or more expressions, then $p(x)$ is called the Common Multiple of the given expressions. The Least Common Multiple (L.C.M) is the product of common factors together with non-common factors of the given expressions.

Finding H.C.F

We can find H.C.F of given expressions by the following two methods.

- (i) By Factorization
- (ii) By division

H.C.F. by Factorization**Example**

Find the H.C.F of the following polynomials.

$$x^2 - 4, x^2 + 4x + 4, 2x^2 + x - 6$$

Solution

By factorization,

$$x^2 - 4 = (x+2)(x-2)$$

$$x^2 + 4x + 4 = (x+2)^2 = (x+2)(x+2)$$

$$\begin{aligned} 2x^2 + x - 6 &= 2x^2 + 4x - 3x - 6 = 2x(x+2) - 3(x+2) \\ &= (x+2)(2x-3) \end{aligned}$$

$$\text{Common factors} = x + 2$$

$$\text{H.C.F} = x + 2$$

H.C.F. by Division**Example**

Use division method to find the H.C.F. of the polynomials

$$p(x) = x^3 - 7x^2 + 14x - 8 \text{ and}$$

$$q(x) = x^3 - 7x + 6$$

Solution

$$\begin{array}{r} 1 \\ \hline x^3 - 7x + 6 \quad | \quad x^3 - 7x^2 + 14x - 8 \\ \quad + x^3 \quad \quad \quad - 7x \quad + 6 \\ \hline \quad \quad \quad \quad \quad + \quad \quad \quad - \\ \quad \quad \quad \quad \quad - 7x^2 + 21x - 14 \end{array}$$

Here the remainder can be factorized as

$$-7x^2 + 21x - 14 = -7(x^2 - 3x + 2)$$

We ignore -7 because it is not common to both the given polynomials and consider $x^2 - 3x + 2$.

$$\begin{array}{r} x+3 \\ \hline x^2 - 3x + 2 \quad | \quad x^3 + 0x^2 - 7x + 6 \\ \quad + x^3 - 3x^2 + 2x \\ \hline \quad \quad \quad \quad \quad - + - \\ \quad \quad \quad \quad \quad 3x^2 - 9x + 6 \\ \quad \quad \quad \quad \quad + \quad - \\ \hline \quad \quad \quad \quad \quad 0 \end{array}$$

Hence H.C. F of $p(x)$ and $q(x)$ is

$$x^2 - 3x + 2$$

Example

Find the L.C.M of $p(x) = 12(x^3 - y^3)$ and $q(x) = 8(x^3 - xy^2)$

Solution

By prime factorization of the given expressions, we have

$$p(x) = 12(x^3 - y^3) = 2^2 \times 3 \times (x-y)(x^2 + xy + y^2) \text{ and}$$

$$q(x) = 8(x^3 - xy^2) = 8x(x^2 - y^2) = 2^3 x(x+y)(x-y) \text{ Hence L.C.M. of } p(x) \text{ and } q(x), \\ 2^3 \times 3 \times x(x+y)(x-y)(x^2 + xy + y^2) = 24x(x+y)(x^3 - y^3)$$

Relation between H.C.F and L.C.M**Example**

By factorization, find (i) H.C.F (ii) L.C.M of $p(x) = 12(x^5 - x^4)$ and $q(x) = 8(x^4 - 3x^3 + 3x^2)$. Establish a relation between $p(x)$, $q(x)$ and H.C.F and L.C.M of the expressions $p(x)$ and $q(x)$.

Solution

Firstly, let us factorize completely the given expressions $p(x)$ and $q(x)$ into irreducible factors. We have

$$p(x) = 12(x^5 - x^4) = 12x^4(x-1) = 2^2 \times 3 \times x^4(x-1) \text{ and}$$

$$q(x) = 8(x^4 - 3x^3 + 2x^2) = 8x^2(x^2 - 3x + 2) = 2^3 x^2(x-1)(x-2)$$

$$\text{H.C.F. of } p(x) \text{ and } q(x) = 2^2 x^2(x-1) = 4x^2(x-1)$$

$$\text{L.C.M of } p(x) \text{ and } q(x) = 2^3 \times 3 \times x^4(x-1)(x-2)$$

$$\begin{aligned} \text{Now } p(x) q(x) &= 12x^4(x-1) \times 8x^2(x-1)(x-2) \\ &= 96x^6(x-1)^2(x-2) \dots \dots \dots \text{(i)} \end{aligned}$$

and (L.C.M) (H.C.F)

$$= [2^3 \times 3 \times x^4(x-1)(x-2)] [4x^2(x-1)]$$

$$= [24x^4(x-1)(x-2)] [4x^2(x-1)]$$

$$= 96x^4(x-1)^2(x-2) \dots \dots \text{(ii)}$$

From (i) and (ii)

$$\text{L.C.M} \times \text{H.C.F} = P(x) \times q(x)$$

Note

$$(1) \quad \text{L.C.M} = \frac{p(x) \times q(x)}{\text{H.C.F}} \quad \text{or}$$

$$\text{H.C.F} = \frac{p(x) \times q(x)}{\text{L.C.M}}$$

(2) If L.C.M, H.C.F and one of $p(x)$ or $q(x)$ are known, then

$$p(x) = \frac{\text{L.C.M} \times \text{H.C.F}}{q(x)}$$

$$q(x) = \frac{\text{L.C.M} \times \text{H.C.F}}{p(x)}$$

Example

Find H.C.F of the polynomials,

$$p(x) = 20(2x^3 + 3x^2 - 2x)$$

$$q(x) = 9(5x^4 + 40x)$$

Then using the above formula (I) find the L.C.M of $p(x)$ and $q(x)$.

Solution

We have

$$p(x) = 20(2x^3 + 3x^2 - 2x) = 20x(2x^2 + 3x - 2)$$

$$= 20x(2x^2 + 4x - x - 2) = 20x[2x(x+2) - (x+2)] = 20x(x+2)(2x-1) = 2^2 \times 5 \times x(x+2)(2x-1)$$

$$q(x) = 9(5x^4 + 40x) = 9x(5x^3 + 40) = 9x[(x^3) + (2)^3]$$

$$= 9x(x+2)(x^2 - 2x + 4) = 9 \times 3^2 \times x(x+2)(x^2 - 2x + 4) \text{ Thus H.C.F of } p(x) \text{ and}$$

$q(x)$ is:

$$= 5x(x+2)$$

$$\text{Now, using the formula} \quad \text{L.C.M.} = \frac{p(x) \times q(x)}{\text{H.C.F}}$$

We obtain

$$\begin{aligned} \text{L.C.M.} &= \frac{2^2 \times 5 \times x(x+2)(2x-1) \times 5 \times 3^2 \times x(x+2)(x^2 - 2x + 4)}{5x(x+2)} \\ &= 4 \times 5 \times 9 \times x(x+2)(2x-1)(x^2 - 2x + 4) \\ &= 180x(x+2)(2x-1)(x^2 - 2x + 4) \end{aligned}$$

Example

Find the L.C.M of

$$p(x) = 6x^3 - 7x^2 - 27x + 8 \quad \text{and}$$

$$q(x) = 6x^3 + 17x^2 + 9x - 4$$

Solution

We have, by long division,

$$\begin{array}{r} 1 \\ \hline 6x^3 - 7x^2 - 27x + 8 \Big| 6x^3 + 17x^2 + 9x - 4 \\ 6x^3 - 7x^2 - 27x + 8 \\ \hline - + + - \\ \hline 24x^2 + 36x - 12 \end{array}$$

But the remainder $24x^2 + 36x - 12$

$$= 12(2x^2 + 3x - 1)$$

Thus, ignoring 12, we have

$$\begin{array}{r} 3x-8 \\ \hline 2x^2 + 3x - 1 \Big| 6x^3 - 7x^2 - 27x + 8 \\ 6x^3 + 9x^2 - 3x \\ \hline - - + \\ \hline -16x^2 - 24x + 8 \\ -16x^2 - 24x + 8 \\ \hline + + - \\ \hline 0 \end{array}$$

Hence H.C.F of $p(x)$ and $q(x)$ is

$$= 2x^2 + 3x - 1$$

$$\begin{array}{r} x^2 - 3x + 2 \\ \underline{x^2 - 3x + 2} \\ \cancel{x^4} + x^2 - 10x + 8 \\ -\cancel{x^4} \pm 3x^2 \pm 2x \\ \hline 4x^2 - 12x + 8 \\ -4x^2 - 12x \pm 8 \\ \hline 0 \end{array}$$

Hence HCF = $x^2 - 3x + 2$

ii) $P(x) = x^4 + x^3 - 2x^2 + x - 3,$
 $q(x) = 5x^3 + 3x^2 - 17x + 6$

$$\begin{array}{r} x^2 + 2 \\ \underline{5x^3 + 3x^2 - 17x + 6} \\ \cancel{5x^4} + 5x^3 - 10x^2 + 5x - 15 \\ -\cancel{5x^4} \pm 3x^3 \mp 17x^2 \pm 6x \\ \hline 2x^3 + 7x^2 - x - 15 \\ \times 5 \\ \hline 10x^4 + 35x^3 - 5x - 75 \\ -10x^4 \pm 6x^3 \mp 34x \pm 12 \\ \hline 29x^2 + 29x - 87 \end{array} \quad (\text{Multiplying by 5})$$

Divided by 29

$$x^2 + x - 3$$

$$\begin{array}{r} 5x - 2 \\ \underline{x^2 + x - 3} \\ \cancel{5x^4} + 3x^2 - 17x + 6 \\ -\cancel{5x^4} \pm 5x^2 \mp 15x \\ \hline 2x^2 - 2x + 6 \\ \mp 2x^2 \mp 2x \pm 6 \\ \hline 0 \end{array}$$

Hence H.C.F = $x^2 + x - 3$

iii) $p(x) = 2x^5 - 4x^4 - 6x,$
 $q(x) = x^5 + x^4 - 3x^3 - 3x^2$

$$\begin{array}{r} 2 \\ \underline{x^5 + x^4 - 3x^3 - 3x^2} \\ \cancel{2x^6} - 4x^4 - 6x \\ -\cancel{2x^6} \pm 2x^4 \mp 6x^3 \mp 6x^2 \\ \hline -6x^4 + 6x^3 + 6x^2 - 6x \end{array}$$

Dividing by -6

$$\begin{array}{r} x^4 - x^3 - x^2 + x \\ \underline{x^4 - x^3 - x^2 + x} \\ \cancel{x^8} + x^4 - 3x^3 - 3x^2 \\ -\cancel{x^8} \mp x^4 \mp x^3 \pm x^2 \\ \hline 2x^4 - 2x^3 - 4x^2 \\ \mp 2x^4 \mp 2x^3 \mp 2x^2 \pm 2x \\ \hline -2x^2 - 2x \end{array}$$

Dividing by -2

$$\begin{array}{r} x^2 - 2x + 1 \\ \underline{x^2 + x} \\ \cancel{x^4} - x^3 - x^2 + x \\ -\cancel{x^4} \pm x^3 \\ \hline 2x^3 - x^2 + x \\ \mp 2x^3 \mp 2x^2 \\ \hline x^2 + x \\ \pm x^2 \pm x \\ \hline 0 \end{array}$$

Hence H.C.F = $x^2 + x = x(x+1)$

Q4. Find the L.C.M of the following expressions:

i) $39x^7y^3z$ and $91x^5y^6z^7$

Sol: By factorization

$$39x^7y^3z = 13 \times 3 \cdot x \cdot x \cdot x \cdot x \cdot x \cdot y \cdot y \cdot y \cdot z$$

$$91x^5y^6z^7 = 13 \times 7 \cdot x \cdot x \cdot x \cdot x \cdot y \cdot y \cdot y \cdot y \cdot y \cdot z \cdot z \cdot z \cdot z \cdot z \cdot z$$

Hence L.C.M =

$$13 \times 3 \times 7 \cdot x \cdot x \cdot x \cdot x \cdot x \cdot y \cdot y \cdot y \cdot y \cdot y \cdot z \cdot z \cdot z \cdot z \cdot z \\ = 273x^7y^6z^7$$

ii) $102xy^2z, 85x^2yz$ and $187xyz^2$

Sol: By factorization

$$102xy^2z = 2 \times 3 \times 17 \cdot x \cdot y \cdot y \cdot z$$

$$85x^2yz = 5 \times 17 \cdot x \cdot x \cdot y \cdot z$$

$$187xyz^2 = 11 \times 17 \cdot x \cdot y \cdot z \cdot z$$

$$\begin{aligned}\text{Hence L.C.M} &= 17 \times 11 \times 5 \times 3 \times 2 \cdot x \cdot x \cdot y \cdot y \cdot z \cdot z \\ &= 5610x^2y^2z^2\end{aligned}$$

Q5. Find the L.C.M of the following expressions by factorization:

i) $x^2 - 25x + 100$ and $x^2 - x - 20$

Sol: By factorization

$$\begin{aligned}x^2 - 25x + 100 &= x^2 - 5x - 20x + 100 \\ &= x(x-5) - 20(x-5) \\ &= (x-5)(x-20) \quad \dots \text{(i)} \\ x^2 - x - 20 &= x^2 - 5x + 4x - 20 \\ &= x(x-5) + 4(x-5) \\ &= (x-5)(x+4) \quad \dots \text{(ii)}\end{aligned}$$

From (i) and (ii)

$$\text{L.C.M} = (x-5)(x-20)(x+4)$$

ii) $x^2 + 4x + 4, x^2 - 4, 2x^2 + x - 6$

Sol: By factorization

$$\begin{aligned}x^2 + 4x + 4 &= x^2 + 2x + 2x + 4 \\ &= x(x+2) + 2(x+2) \\ &= (x+2)(x+2) \quad \dots \text{(i)} \\ x^2 - 4 &= (x)^2 - (2)^2 \\ &= (x+2)(x-2) \quad \dots \text{(ii)}\end{aligned}$$

$$\begin{aligned}2x^2 + x - 6 &= 2x^2 + 4x - 3x - 6 \\ &= 2x(x+2) - 3(x+2) \\ &= (x+2)(2x-3) \quad \dots \text{(iii)}\end{aligned}$$

From (i), (ii) and (iii)

$$\begin{aligned}\text{LCM} &= (x+2)(x+2)(x-2)(2x-3) \\ &= (x+2)^2(x-2)(2x-3)\end{aligned}$$

iii) $2(x^4 - y^4), 3(x^3 + 2x^2y - xy^2 - 2y^3)$

Sol: By factorization

$$2(x^4 - y^4) = 2[(x^2)^2 - (y^2)^2]$$

$$\begin{aligned}&= 2(x^2 + y^2)(x^2 - y^2) \\ &= 2(x^2 + y^2)(x+y)(x-y) \quad \dots \text{(i)}\end{aligned}$$

$$\begin{aligned}3(x^3 + 2x^2y - xy^2 - 2y^3) &= 3[x^2(x+2y) - y^2(x+2y)] \\ &= 3(x+2y)(x^2 - y^2) \\ &= 3(x+2y)(x+y)(x-y) \quad \dots \text{(ii)}\end{aligned}$$

From (i) & (ii)

L.C.M =

$$\begin{aligned}&2 \times 3(x+y)(x-y)(x^2 + y^2)(x+2y) \\ &= 6(x^4 - y^4)(x+2y)\end{aligned}$$

iv) $4(x^4 - 1), 6(x^3 - x^2 - x + 1)$

Sol: By factorization

$$\begin{aligned}4(x^4 - 1) &= 4[(x^2)^2 - (1)^2] \\ &= 4(x^2 + 1)(x^2 - 1) \\ &= 2 \times 2(x^2 + 1)[(x)^2 - (1)^2] \\ &= 2 \times 2(x^2 + 1)(x+1)(x-1) \quad \dots \text{(i)} \\ 6(x^3 - x^2 - x + 1) &= 6[x^2(x-1) - 1(x-1)] \\ &= 6(x-1)(x^2 - 1) = 2 \times 3(x-1)[(x)^2 - (1)^2] \\ &= 2 \times 3(x-1)(x-1)(x+1) \quad \dots \text{(ii)}\end{aligned}$$

From (i) & (ii)

$$\begin{aligned}\text{LCM} &= 2 \times 2 \times 3(x+1)(x-1)(x^2 + 1)(x-1) \\ &= 12(x^4 - 1)(x-1)\end{aligned}$$

Q6. For what value of k is $(x+4)$, the H.C.F of $x^2 + x - (2k+2)$ and $2x^2 + kx - 12$?

Sol: $k = ?$

$$\begin{aligned}p(x) &= x^2 + x - (2k+2) \text{ and} \\ q(x) &= 2x^2 + kx - 12\end{aligned}$$

As given that $x+4$ is HCF, so $p(x)$ and $q(x)$ will be exactly divisible by $(x+4)$

$$\begin{array}{r}
 \frac{x-3}{x+4} \\
 \overline{x^2 + x - (2k+2)} \\
 \underline{-x^2 - 4x} \\
 \overline{-3x - (2k+2)} \\
 \underline{-3x - 12} \\
 \overline{12 - (2k+2)} \\
 = 12 - 2k - 2 \\
 = 10 - 2k
 \end{array}$$

As $p(x)$ is exactly divisible by $x+4$, so,

$$10 - 2k = 0$$

$$10 = 2k$$

$$\frac{10}{2} = k$$

$$k = 5$$

Q7. If $(x+3)(x-2)$ is the H.C.F of $p(x) = (x+3)(2x^2 - 3x + k)$ and $q(x) = (x-2)(3x^2 + 7x - l)$, find k and l .

Sol: $k = ?$ and $l = ?$

As $(x+3)(x-2)$ is the H.C.F, so $p(x)$ and $q(x)$ will be exactly divisible by

$(x+3)(x-2)$ i.e., $\frac{p(x)}{\text{HCF}}$ has remainder zero.

$$\frac{(x+3)(2x^2 - 3x + k)}{(x+3)(x-2)} = \frac{2x^2 - 3x + k}{x-2}$$

$$\begin{array}{r}
 \frac{2x+1}{x-2} \\
 \overline{2x^2 - 3x + k} \\
 \underline{-2x^2 + 4x} \\
 \overline{x+k} \\
 \underline{-x-2} \\
 \overline{k+2}
 \end{array}$$

As remainder = 0, then

$$k+2=0$$

$$k = -2$$

and $\frac{q(x)}{\text{HCF}}$ has zero remainder

$$\frac{(x-2)(3x^2 + 7x - l)}{(x+3)(x-2)} = \frac{3x^2 + 7x - l}{x+3}$$

$$\begin{array}{r}
 \frac{3x-2}{x+3} \\
 \overline{3x^2 + 7x - l} \\
 \underline{-3x^2 - 9x} \\
 \overline{2x - l} \\
 \underline{-2x - 6} \\
 \overline{-l + 6}
 \end{array}$$

As remainder = 0

$$-l+6=0$$

$$-l=-6$$

$$\Rightarrow l=6$$

Q8. The LCM and HCF of two polynomials $p(x)$ and $q(x)$ are $2(x^4 - 1)$ and $(x+1)(x^2 + 1)$ respectively. If $p(x) = x^3 + x + 1$, find $q(x)$.

Sol: LCM = $2(x^4 - 1)$,

$$\text{HCF} = (x+1)(x^2 + 1)$$

$$p(x) = x^3 + x^2 + x + 1, q(x) = ?$$

As $p(x) \times q(x) = (\text{LCM}) \times (\text{HCF})$

$$q(x) = \frac{(\text{LCM}) \times (\text{HCF})}{p(x)}$$

$$= \frac{2(x^4 - 1) \times (x+1)(x^2 + 1)}{x^3 + x^2 + x + 1}$$

$$= \frac{2(x^4 - 1)(x^3 + x^2 + x + 1)}{x^3 + x^2 + x + 1}$$

$$q(x) = 2(x^4 - 1)$$

Q9. Let $p(x) = 10(x^2 - 9)(x^2 - 3x + 2)$ and $q(x) = 10x(x+3)(x-1)^2$. If the H.C.F. of $p(x), q(x)$ is $10(x+3)(x-1)$, find their L.C.M.

Sol: $p(x) = 10(x^2 - 9)(x^2 - 3x + 2)$,

$$q(x) = 10x(x+3)(x-1)^2$$

$$\text{H.C.F.} = 10(x+3)(x-1), \text{ L.C.M.} = ?$$

$$\text{As } (\text{L.C.M.}) \times (\text{H.C.F.}) = p(x) \times q(x)$$

$$\text{L.C.M.} = \frac{p(x) \times q(x)}{\text{H.C.F.}}$$

$$= \frac{10(x^2 - 9)(x^2 - 3x + 2) \times 10x(x+3)(x-1)^2}{10(x+3)(x-1)}$$

$$= \frac{(x^2 - 9)(x^2 - 3x + 2) \times 10x \cancel{(x+3)} \cancel{(x-1)} (x-1)}{\cancel{(x+3)} \cancel{(x-1)}}$$

$$= 10x(x-1)(x^2 - 9)(x^2 - 3x + 2)$$

$$= 10x(x-1)(x^2 - 9)(x^2 - x - 2x + 2)$$

$$= 10x(x-1)(x^2 - 9)[x(x-1) - 2(x-1)]$$

$$= 10x(x-1)(x^2 - 9)(x-1)(x-2)$$

$$= 10x(x-1)^2(x^2 - 9)(x-2)$$

Q10. Let the product of L.C.M and H.C.F of two polynomials be $(x+3)^2(x-2)(x+5)$. If one polynomial is $(x+3)(x-2)$ and the second polynomial is $x^2 + kx + 15$, find the value of k .

Sol: $k = ?$

Product of L.C.M. & H.C.F is

$$\text{LCM} \times \text{HCF} = (x+3)^2(x-2)(x+5)$$

$$p(x) = (x+3)(x-2)$$

$$q(x) = x^2 + kx + 15$$

$$\text{As } p(x) \times q(x) = \text{LCM} \times \text{HCF}$$

$$(x+3)(x-2)(x^2 + kx + 15)$$

$$= (x+3)^2(x-2)(x+5)$$

$$x^2 + kx + 15 = \frac{(x+3)(x+3)(x-2)(x+5)}{(x+3)(x-2)}$$

$$x^2 + kx + 15 = (x+3)(x+5)$$

$$x^2 + kx + 15 = x^2 + 3x + 5x + 15$$

$$x^2 + kx + 15 = x^2 + 8x + 15$$

Comparing co-efficient of ' x '

$$\Rightarrow kx = 8x$$

$$[k = 8]$$

Q11. Waqas wishes to distribute 128 bananas and also 176 apples equally among a certain number of children. Find the highest number of the Children. Who can get the fruit in this way?

Sol: No. of bananas = 128

No. of apples = 176

Highest no. of children who get the fruit in this way is H.C.F.

So

No. of bananas =

$$2 \times 2 \times 2 \times 2 \times 2 \times 2$$

No. of apples =

$$2 \times 2 \times 2 \times 11$$

Hence required no. of children =

$$2 \times 2 \times 2 \times 2 = 16$$

Example

Simplify

$$\frac{x+3}{x^2 - 3x + 2} + \frac{x+2}{x^2 - 4x + 3} + \frac{x+1}{x^2 - 5x + 6}, x \neq 1, 2, 3$$

Solution

$$\frac{x+3}{x^2 - 3x + 2} + \frac{x+2}{x^2 - 4x + 3} + \frac{x+1}{x^2 - 5x + 6}$$

$$= \frac{x+3}{x^2 - 2x - x + 2} + \frac{x+2}{x^2 - 3x - x + 3} + \frac{x+1}{x^2 - 3x - 2x + 6}$$

$$= \frac{x+3}{x(x-2) - 1(x-2)} + \frac{x+2}{x(x-3) - 1(x-3)} + \frac{x+1}{x(x-3) - 2(x-3)}$$

$$\begin{aligned}
 &= \frac{x+3}{(x-2)(x-1)} + \frac{x+2}{(x-3)(x-1)} + \frac{x+1}{(x-3)(x-2)} \\
 &= \frac{(x+3)(x-3)+(x+2)(x-2)+(x+1)(x-1)}{(x-1)(x-2)(x-3)} \\
 &= \frac{x^2-9+x^2-4+x^2-1}{(x-1)(x-2)(x-3)} \\
 &= \frac{3x^2-14}{(x-1)(x-2)(x-3)}
 \end{aligned}$$

Example

$$\text{Express the product } \frac{x^3-8}{x^2-4} \times \frac{x^2+6x+8}{x^2-2x+1}$$

as an algebraic expression reduced lowest forms $x \neq 2, -2, 1$

Solution

By factorizing completely, we have

$$\begin{aligned}
 &\frac{x^3-8}{x^2-4} \times \frac{x^2+6x+8}{x^2-2x+1} \\
 &= \frac{(x-2)(x^2+2x+4) \times (x+2)(x+4)}{(x-2)(x+2) \times (x-1)^2} \dots(i)
 \end{aligned}$$

Now the factors of numerator are $(x-2), (x^2+2x+4), (x+2)$ and $(x+4)$ and the factors of denominator are

$(x-2), (x+2)$ and $(x-1)^2$.

Therefore, their H.C.F. is $(x-2) \times (x+2)$

By cancelling H.C.F i.e., $(x-2) \times (x+2)$ from (i), we get the simplified form of given product as the fraction $\frac{(x^2+2x+4)(x+4)}{(x-1)^2}$

Example

$$\text{Divide } \frac{x^2+x+1}{x^2-9} \text{ by } \frac{x^3-1}{x^2-4x+3}$$

and simplify by reducing to lowest forms.

Solution

$$\begin{aligned}
 &\text{We have } \frac{x^2+x+1}{x^2-9} \div \frac{x^3-1}{x^2-4x+3} \\
 &= \frac{(x^2+x+1)}{(x^2-9)} \times \frac{(x^2-4x+3)}{(x^3-1)} \\
 &= \frac{(x^2+x+1)(x^2-x-3x+3)}{(x^2-9)(x^3-1)} \\
 &= \frac{\left(x^2+x+1\right)\left[x(x-1)-3(x-1)\right]}{(x+3)(x-3)(x-1)(x^2+x+1)} \\
 &= \frac{(x^2+x+1)(x-3)(x-1)}{(x+3)(x-3)(x-1)(x^2+x+1)} = \frac{1}{x+3}, x \neq -3
 \end{aligned}$$

Exercise 6.2

Simplify each of the following as a rational expression.

$$\begin{aligned}
 \text{Q1. } &\frac{x^2-x-6}{x^2-9} + \frac{x^2+2x-24}{x^2-x-12} \\
 &= \frac{x^2-3x+2x-6}{(x-3)^2} + \frac{x^2+6x-4x-24}{x^2+3x-4x-12}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{x(x-3)+2(x-3)}{(x+3)(x-3)} + \frac{x(x+6)-4(x+6)}{x(x+3)-4(x+3)} \\
 &= \frac{(x-3)(x+2)}{(x+3)(x-3)} + \frac{(x+6)(x-4)}{(x+3)(x-4)} \\
 &= \frac{x+2}{x+3} + \frac{x+6}{x+3} = \frac{x+2+x+6}{x+3}
 \end{aligned}$$

$$= \frac{2x+8}{x+3}$$

$$= \frac{2(x+4)}{x+3}$$

Q2.

$$\begin{aligned} & \left[\frac{x+1}{x-1} - \frac{x-1}{x+1} - \frac{4x}{x^2+1} \right] + \frac{4x}{x^4-1} \\ & = \left[\frac{(x+1)^2 - (x-1)^2}{(x-1)(x+1)} - \frac{4x}{x^2+1} \right] + \frac{4x}{x^4-1} \\ & = \left[\frac{(x^2+2x+1) - (x^2-2x+1)}{(x)^2-(1)^2} - \frac{4x}{x^2+1} \right] + \frac{4x}{x^4-1} \\ & = \left[\frac{x^2+2x+1-x^2+2x-1}{x^2-1} - \frac{4x}{x^2+1} \right] + \frac{4x}{x^4-1} \\ & = \left[\frac{4x}{x^2-1} - \frac{4x}{x^2+1} \right] + \frac{4x}{x^4-1} \\ & = \left[\frac{4x(x^2+1) - 4x(x^2-1)}{(x^2-1)(x^2+1)} \right] + \frac{4x}{x^4-1} \\ & = \frac{4x^3+4x-4x^3+4x}{(x^2)^2-(1)^2} + \frac{4x}{x^4-1} \end{aligned}$$

$$= \frac{8x}{x^4-1} + \frac{4x}{x^4-1}$$

$$= \frac{8x+4x}{x^4-1}$$

$$= \frac{12x}{x^4-1}$$

Q3.

$$\begin{aligned} & \frac{1}{x^2-8x+15} + \frac{1}{x^2-4x+3} - \frac{2}{x^2-6x+5} \\ & = \frac{1}{x^2-3x-5x+15} + \frac{1}{x^2-3x-x+3} - \frac{2}{x^2-5x-x+5} \\ & = \frac{1}{x(x-3)-5(x-3)} + \frac{1}{x(x-3)-1(x-3)} - \frac{2}{x(x-5)-1(x-5)} \end{aligned}$$

$$= \frac{1}{(x-3)(x-5)} + \frac{1}{(x-3)(x-1)} - \frac{2}{(x-5)(x-1)}$$

$$= \frac{x-1+x-5-2(x-3)}{(x-1)(x-3)(x-5)}$$

$$= \frac{x-1+x-5-2x+6}{(x-1)(x-3)(x-5)}$$

$$= \frac{2x-6-2x+6}{(x-1)(x-3)(x-5)}$$

$$= \frac{0}{(x-1)(x-3)(x-5)}$$

$$= 0$$

Q4.

$$\frac{(x+2)(x+3)}{x^2-9} + \frac{(x+2)(2x^2-32)}{(x-4)(x^2-x-6)}$$

$$= \frac{(x+2)(x+3)}{(x)^2-(3)^2} + \frac{(x+2).2(x^2-16)}{(x-4)(x^2+2x-3x-6)}$$

$$= \frac{(x+2)(x+3)}{(x-3)(x+3)} + \frac{2(x+2)[(x)^2-(4)^2]}{(x-4)(x^2+2x-3x-6)}$$

$$= \frac{(x+2)}{x-3} + \frac{2(x+2)(x+4)(x-4)}{(x-4)(x+2)(x-3)}$$

$$= \frac{x+2}{x-3} + \frac{2x+8}{x-3}$$

$$= \frac{x+2+2x+8}{x-3}$$

$$= \frac{3x+10}{x-3}$$

Q5.

$$\frac{x+3}{2x^2+9x+9} + \frac{1}{2(2x-3)} - \frac{4x}{4x^2-9}$$

$$= \frac{x+3}{2x^2+6x+3x+9} + \frac{1}{2(2x-3)} - \frac{4x}{(2x)^2-(3)^2}$$

$$= \frac{x+3}{2x(x+3)+3(x+3)} + \frac{1}{2(2x-3)} - \frac{4x}{(2x+3)(2x-3)}$$

$$\begin{aligned}
&= \frac{x+3}{(x+3)(2x+3)} + \frac{1}{2(2x-3)} - \frac{4x}{(2x+3)(2x-3)} \\
&= \frac{1}{2x+3} + \frac{1}{2(2x-3)} - \frac{4x}{(2x+3)(2x-3)} \\
&= \frac{2(2x-3) + 2x+3 - 2(4x)}{2(2x+3)(2x-3)} \\
&= \frac{4x-6+2x+3-8x}{2(2x+3)(2x-3)} \\
&= \frac{-2x-3}{2(2x+3)(2x-3)} \\
&= \frac{-1(2x+3)}{2(2x+3)(2x-3)} \\
&= \frac{-1}{2(2x-3)} \\
&= \frac{1}{2(3-2x)}
\end{aligned}$$

Q6. $A - \frac{1}{A}$, where $A = \frac{a+1}{a-1}$

so $\frac{1}{A} = \frac{a-1}{a+1}$

$$\begin{aligned}
\text{Now } A - \frac{1}{A} &= \frac{a+1}{a-1} - \frac{a-1}{a+1} \\
&= \frac{(a+1)^2 - (a-1)^2}{(a-1)(a+1)} \\
&= \frac{(a^2 + 2a + 1) - (a^2 - 2a + 1)}{(a)^2 - (1)^2} \\
&= \frac{a^2 + 2a + 1 - a^2 + 2a - 1}{a^2 - 1} \\
&= \frac{4a}{a^2 - 1}
\end{aligned}$$

$$\begin{aligned}
\text{Q7. } &\left[\frac{x-1}{x-2} + \frac{2}{2-x} \right] - \left[\frac{x+1}{x+2} + \frac{4}{4-x^2} \right] \\
&= \left[\frac{-(x-1)}{2-x} + \frac{2}{2-x} \right] - \left[\frac{x+1}{x+2} + \frac{4}{(2)^2 - (x)^2} \right] \\
&= \left[\frac{-(x-1)}{2-x} + \frac{2}{2-x} \right] - \left[\frac{x+1}{x+2} + \frac{4}{(2+x)(2-x)} \right] \\
&= \left[\frac{-x+1+2}{2-x} \right] - \left[\frac{(x+1)(2-x)+4}{(2+x)(2-x)} \right] \\
&= \frac{3-x}{2-x} - \left[\frac{2x-x^2+2-x+4}{(2+x)(2-x)} \right] \\
&= \frac{3-x}{2-x} - \left[\frac{6+x-x^2}{(2+x)(2-x)} \right] \\
&= \frac{3-x}{2-x} - \left[\frac{6+3x-2x-x^2}{(2+x)(2-x)} \right] \\
&= \frac{3-x}{2-x} - \left[\frac{3(2+x)-x(2+x)}{(2+x)(2-x)} \right] \\
&= \frac{3-x}{2-x} - \left[\frac{(2+x)(3-x)}{(2+x)(2-x)} \right] \\
&= \frac{3-x}{2-x} - \frac{3-x}{2-x} \\
&= \frac{3-x-3+x}{2-x} \\
&= \frac{0}{2-x} \\
&= 0
\end{aligned}$$

Q8. What rational expression should be subtracted from $\frac{2x^2 + 2x - 7}{x^2 + x - 6}$ to get

$$\frac{x-1}{x-2} = ?$$

Sol: Let the required expression be A,
then $\frac{2x^2+2x-7}{x^2+x-6} - A = \frac{x-1}{x-2}$

or $\frac{2x^2+2x-7}{x^2+x-6} - \frac{x-1}{x-2} = A$

So $A = \frac{2x^2+2x-7}{x^2+3x-2x-6} - \frac{x-1}{x-2}$
 $= \frac{2x^2+2x-7}{x(x+3)-2(x+3)} - \frac{x-1}{x-2}$
 $= \frac{2x^2+2x-7}{(x+3)(x-2)} - \frac{x-1}{x-2}$
 $= \frac{2x^2+2x-7-(x-1)(x+3)}{(x+3)(x-2)}$
 $= \frac{2x^2+2x-7-(x^2-x+3x-3)}{(x+3)(x-2)}$
 $= \frac{(2x^2+2x-7)-(x^2+2x-3)}{(x+3)(x-2)}$
 $= \frac{2x^2+2x-7-x^2-2x+3}{(x+3)(x-2)}$
 $= \frac{x^2-4}{(x+3)(x-2)}$
 $= \frac{(x)^2-(2)^2}{(x+3)(x-2)}$
 $= \frac{(x+2)(x-2)}{(x+3)(x-2)}$
 $= \frac{x+2}{x+3}$

Perform the indicated operations and simplify to the lowest forms.

Q9. $\frac{x^2+x-6}{x^2-x-6} \times \frac{x^2-4}{x^2-9}$

$$\begin{aligned}
&= \frac{x^2+3x-2x-6}{x^2-3x+2x-6} \times \frac{(x)^2-(2)^2}{(x)^2-(3)^2} \\
&= \frac{x(x+3)-2(x+3)}{x(x-3)+2(x-3)} \times \frac{(x+2)(x-2)}{(x+3)(x-3)} \\
&= \frac{\cancel{(x+3)}(x-2)}{\cancel{(x-3)}(\cancel{x+2})} \times \frac{\cancel{(x+2)}(x-2)}{\cancel{(x+3)}(\cancel{x-3})} \\
&= \frac{(x-2)^2}{(x-3)^2} \\
\textbf{Q10.} \quad & \frac{x^3-8}{x^2-4} \times \frac{x^2+6x+8}{x^2-2x+1} \\
&= \frac{(x)^3-(2)^3}{(x)^2-(2)^2} \times \frac{x^2+2x+4x+8}{x^2-x-x+1} \\
&= \frac{\cancel{(x-2)}[(x)^2+(x)(2)+(2)^2]}{\cancel{(x-2)}(x+2)} \times \frac{x(x+2)+4(x+2)}{x(x-1)-1(x-1)} \\
&= \frac{x^2+2x+4}{\cancel{x+2}} \times \frac{\cancel{(x+2)}(x+4)}{\cancel{(x-1)}(x-1)} \\
&= \frac{(x^2+2x+4)(x+4)}{(x-1)^2} \\
\textbf{Q11.} \quad & \frac{x^4-8x}{2x^2+5x-3} \times \frac{2x-1}{x^2+2x+4} \times \frac{x+3}{x^2-2x} \\
&= \frac{x(x^3-8)}{2x^2+6x-x-3} \times \frac{2x-1}{x^2+2x+4} \times \frac{x+3}{x(x-2)} \\
&= \frac{x[(x)^3-(2)^3]}{2x(x+3)-1(x+3)} \times \frac{2x-1}{x^2+2x+4} \times \frac{x+3}{x(x-2)} \\
&= \frac{\cancel{x}(x-2)(x^2+2x+4)}{\cancel{(x+3)}(2x-1)} \times \frac{2x-1}{\cancel{x^2+2x+4}} \times \frac{x+3}{\cancel{x}(x-2)} \\
&= 1
\end{aligned}$$

Q12. $\frac{2y^2+7y-4}{3y^2-13y+4} \div \frac{4y^2-1}{6y^2+y-1}$

$$\begin{aligned}
&= \frac{2y^2 + 8y - y - 4}{3y^2 - y - 12y + 4} \div \frac{(2y)^2 - (1)^2}{6y^2 + 3y - 2y - 1} \\
&= \frac{2y(y+4) - 1(y+4)}{y(3y-1) - 4(3y-1)} \div \frac{(2y+1)(2y-1)}{3y(2y+1) - 1(2y+1)} \\
&= \frac{(y+4)(2y-1)}{(3y-1)(y-4)} \div \frac{(2y+1)(2y-1)}{(2y+1)(3y-1)} \\
&= \frac{(y+4)(2y-1)}{(3y-1)(y-4)} \times \frac{(2y+1)(3y-1)}{(2y+1)(2y-1)} \\
&= \frac{y+4}{y-4}
\end{aligned}$$

Q13. $\left[\frac{x^2+y^2}{x^2-y^2} - \frac{x^2-y^2}{x^2+y^2} \right] \div \left[\frac{x+y}{x-y} - \frac{x-y}{x+y} \right]$

$$\begin{aligned}
&= \left[\frac{(x^2+y^2)^2 - (x^2-y^2)^2}{(x^2-y^2)(x^2+y^2)} \right] \div \left[\frac{(x+y)^2 - (x-y)^2}{(x-y)(x+y)} \right] \\
&= \frac{x^4 + y^4 + 2x^2y^2 - (x^4 + y^4 - 2x^2y^2)}{(x^2-y^2)(x^2+y^2)} \\
&\div \frac{x^2 + y^2 + 2xy - x^2 - y^2 + 2xy}{x^2 - y^2} \\
&= \frac{x^4 + y^4 + 2x^2y^2 - x^4 - y^4 + 2x^2y^2}{(x^2-y^2)(x^2+y^2)} \\
&\div \frac{x^4 + y^4 + 2xy - x^4 - y^4 + 2xy}{x^2 - y^2} \\
&= \frac{4x^2y^2}{(x^2-y^2)(x^2+y^2)} \div \frac{4xy}{x^2-y^2} \\
&= \frac{4x^2y^2}{(x^2-y^2)(x^2+y^2)} \times \frac{x^2-y^2}{4xy} \\
&= \frac{xy}{x^2+y^2}
\end{aligned}$$

Square Root of Algebraic Expression

The square root of a given expression $p(x)$ as another expression $q(x)$ such that $q(x) \cdot q(x) = p(x)$.

As $5 \times 5 = 25$, so square root of 25 is 5

It means we can find square root of the expression $p(x)$ if it can be expressed as a perfect square.

Example

Use factorization to find the square root of the expression

$$4x^2 - 12x + 9$$

Solution

$$\text{We have, } 4x^2 - 12x + 9$$

$$\begin{aligned}
&= 4x^2 - 6x - 6x + 9 = 2x(2x-3) - 3(2x-3) \\
&= (2x-3)(2x-3) = (2x-3)^2
\end{aligned}$$

$$\text{Hence } \sqrt{4x^2 - 12x + 9}$$

$$= \pm(2x-3)$$

Example

Find the square root of $x^2 + \frac{1}{x^2} + 12\left(x + \frac{1}{x}\right) + 38, x \neq 0$

Solution

$$\begin{aligned}
&\text{We have } x^2 + \frac{1}{x^2} + 12\left(x + \frac{1}{x}\right) + 38 \\
&= x^2 + \frac{1}{x^2} + 2 + 12\left(x + \frac{1}{x}\right) + 36, \\
&\quad (\text{adding and subtracting 2})
\end{aligned}$$

$$\begin{aligned}
 &= \left(x + \frac{1}{x} \right)^2 + 2 \left(x + \frac{1}{x} \right) (6) + (6)^2 \\
 &= \left[\pm \left(x + \frac{1}{x} + 6 \right) \right]^2;
 \end{aligned}$$

since $a^2 + 2ab + b^2 = (a+b)^2$

Hence the required square root is
 $\pm \left(x + \frac{1}{x} + 6 \right)$

Example

Find the square root of
 $4x^4 + 12x^3 + x^2 - 12x + 4$

Solution

$$\begin{array}{r}
 2x^2 + 3x - 2 \\
 \hline
 4x^4 + 12x^3 + x^2 - 12x + 4 \\
 4x^4 \\
 \hline
 12x^3 + x^2 - 12x + 4 \\
 12x^3 \pm 9x^2 \\
 \hline
 -8x^2 - 12x + 4 \\
 \mp 8x^2 \mp 12x \pm 4 \\
 \hline
 0
 \end{array}$$

Thus square root of given expression is $\pm(2x^2 + 3x - 2)$

Example 2

Find the square root of the expression

$$4\frac{x^2}{y^2} + 8\frac{x}{y} + 16 + 12\frac{y}{x} + 9\frac{y^2}{x^2}$$

Solution

We note that the given expression is in descending powers of x.

$$\begin{array}{r}
 2\frac{x}{y} + 2 + 3\frac{y}{x} \\
 \hline
 4\frac{x^2}{y^2} + 8\frac{x}{y} + 16 + 12\frac{y}{x} + 9\frac{y^2}{x^2} \\
 4\frac{x^2}{y^2} \\
 \hline
 \cancel{+ 4\frac{x^2}{y^2}} \\
 8\frac{x}{y} + 16 \\
 \pm 8\frac{x}{y} \pm 4 \\
 \hline
 4\frac{x}{y} + 4 + 3\frac{y}{x} \\
 12 + 12\frac{y}{x} + 9\frac{y^2}{x^2} \\
 \pm 12 + 12\frac{y}{x} \pm 9\frac{y^2}{x^2} \\
 \hline
 0
 \end{array}$$

Hence the square root of given expression is $\pm \left(2\frac{x}{y} + 2 + 3\frac{y}{x} \right)$

Example

To make the expression $x^4 - 10x^3 + 33x^2 - 42x + 20$ a perfect square,

- What should be added to it?
- What should be subtracted from it?
- What should be the value of x?

$$\begin{array}{r}
 x^2 - 5x + 4 \\
 \hline
 x^4 - 10x^3 + 33x^2 - 42x + 20 \\
 \pm x^4 \\
 \hline
 -10x^3 + 33x^2 \\
 -10x^3 + 25x^2 \\
 + - \\
 \hline
 8x^2 - 42x + 20 \\
 -8x^2 - 40x + 16 \\
 + - \\
 \hline
 -2x + 4
 \end{array}$$

For making the given expression a perfect square the remainder must be zero.

Hence

- (i) We should add $(2x-4)$ to the given expression
- (ii) We should subtract $(-2x+4)$ from the given expression

(iii) We should take $-2x+4=0$ to find the value of x . This gives the required value of x i.e., $x=2$.

Exercise 6.3

Q1. Use factorization to find the square root of the following expressions.

$$\begin{aligned} \text{i)} \quad & 4x^2 - 12xy + 9y^2 \\ & = (2x)^2 - 2(2x)(3y) + (3y)^2 \\ & = (2x - 3y)^2 \end{aligned}$$

$$\begin{aligned} \text{Hence } & \sqrt{4x^2 - 12xy + 9y^2} \\ & = \sqrt{(2x - 3y)^2} \\ & = \pm(2x - 3y) \end{aligned}$$

$$\begin{aligned} \text{ii)} \quad & x^2 - 1 + \frac{1}{4x^2} \\ & = (x)^2 - 2(x)\left(\frac{1}{2x}\right) + \left(\frac{1}{2x}\right)^2 \end{aligned}$$

$$\begin{aligned} \text{Hence } & \sqrt{x^2 - 1 + \frac{1}{4x^2}} \\ & = \sqrt{\left(x - \frac{1}{2x}\right)^2} \\ & = \pm\left(x - \frac{1}{2x}\right) \end{aligned}$$

$$\begin{aligned} \text{iii)} \quad & \frac{1}{16}x^2 - \frac{1}{12}xy + \frac{1}{36}y^2 \\ & = \left(\frac{1}{4}x\right)^2 - 2\left(\frac{1}{4}x\right)\left(\frac{1}{6}y\right) + \left(\frac{1}{6}y\right)^2 \end{aligned}$$

$$= \left(\frac{1}{4}x - \frac{1}{6}y\right)^2$$

$$\text{Hence } \sqrt{\frac{1}{16}x^2 - \frac{1}{12}xy + \frac{1}{36}y^2}$$

$$= \sqrt{\left(\frac{1}{4}x - \frac{1}{6}y\right)^2}$$

$$= \pm\left(\frac{1}{4}x - \frac{1}{6}y\right)$$

$$\begin{aligned} \text{iv)} \quad & 4(a+b)^2 - 12(a^2 - b^2) + 9(a-b)^2 \\ & = [2(a+b)]^2 - 2 \times 2(a+b) \times 3(a-b) + [3(a-b)]^2 \\ & = [2(a+b) - 3(a-b)]^2 \end{aligned}$$

$$= (-a+5b)^2$$

$$= (5b-a)^2$$

$$\begin{aligned} \text{Hence } & \sqrt{4(a+b)^2 - 12(a^2 - b^2) + 9(a-b)^2} \\ & = \sqrt{(5b-a)^2} \end{aligned}$$

$$= \pm(5b-a)$$

$$\begin{aligned} \text{v)} \quad & \frac{4x^6 - 12x^3y^3 + 9y^6}{9x^4 + 24x^2y^2 + 16y^4} \\ & = \frac{\left(2x^3\right)^2 - 2(2x^3)(3y^3) + (3y^3)^2}{\left(3x^2\right)^2 + 2(3x^2)(4y^2) + (4y^2)^2} \end{aligned}$$

$$= \frac{(2x^3 - 3y^3)^2}{(3x^2 + 4y^2)^2}$$

$$\begin{aligned} \text{Hence } & \sqrt{\frac{4x^6 - 12x^3y^3 + 9y^6}{9x^4 + 24x^2y^2 + 16y^4}} \\ &= \sqrt{\left(\frac{2x^3 - 3y^3}{3x^2 + 4y^2}\right)^2} \\ &= \pm \left(\frac{2x^3 - 3y^3}{3x^2 + 4y^2}\right). \end{aligned}$$

$$\begin{aligned} \text{vi) } & \left(x + \frac{1}{x} \right)^2 - 4 \left(x - \frac{1}{x} \right) \quad (x \neq 0) \\ &= \left(x \right)^2 + \left(\frac{1}{x} \right)^2 + 2 \left(x \right) \left(\frac{1}{x} \right) - 4 \left(x - \frac{1}{x} \right) \\ &= x^2 + \frac{1}{x^2} + 2 - 4 \left(x - \frac{1}{x} \right), \dots \dots \text{(i)} \end{aligned}$$

$$\text{Let } x - \frac{1}{x} = a$$

$$\text{Squaring } \left(x - \frac{1}{x}\right)^2 = (a)^2$$

$$x^2 + \frac{1}{x^2} - 2 = a^2$$

$$x^2 + \frac{1}{r^2} = a^2 + 2$$

So expression (i) becomes

$$= a^2 + 2 + 2 - 4a$$

$$= a^2 - 4a + 4$$

$$= (a)^2 - 2(a)(2) + (2)^2$$

$$= (a - 2)^2$$

Putting value of ' a '

$$= \left(x - \frac{1}{x} - 2 \right)^2$$

$$\text{Hence } = \sqrt{\left(x - \frac{1}{x} - 2\right)^2}$$

$$= \pm \left(x - \frac{1}{x} - 2\right)$$

$$\text{vii) } \left(x^2 + \frac{1}{x^2}\right)^2 - 4\left(x + \frac{1}{x}\right)^2 + 12 \dots \text{(i)}$$

$$\text{Let } x + \frac{1}{x} = a$$

$$\text{Squaring} \quad \left(x + \frac{1}{x} \right)^2 = (a)^2$$

$$x^2 + \frac{1}{x^2} + 2 = a^2$$

$$x^2 + \frac{1}{x^2} = a^2 - 2$$

So expression (i) becomes

$$= (a^2 - 2)^2 - 4(a)^2 + 12$$

$$= (a^2)^2 - 2(a^2)(2) + (2)^2 - 4a^2 + 12$$

$$= a^4 - 4a^2 + 4 - 4a^2 + 12$$

$$= a^4 - 8a^2 + 16$$

$$= (a^2)^2 - 2(a^2)(4) + (4)^2$$

$$= (a^2 - 4)^2$$

Putting values of a^2

$$= \left(x^2 + \frac{1}{x^2} + 2 - 4 \right)^2$$

$$= \left(x^2 + \frac{1}{x^2} - 2 \right)^2$$

$$\text{Hence } = \sqrt{\left(x^2 + \frac{1}{x^2}\right)^2 - 4\left(x + \frac{1}{x}\right)^2 + 12}$$

$$= \sqrt{\left(x^2 + \frac{1}{2} - 2\right)^2}$$

$$= \pm \left(x^2 + \frac{1}{x^2} - 2 \right)$$

viii)
$$\begin{aligned} & (x^2 + 3x + 2)(x^2 + 4x + 3)(x^2 + 5x + 6) \\ & = (x^2 + x + 2x + 2)(x^2 + x + 3x + 3)(x^2 + 2x + 3x + 6) \\ & = [x(x+1) + 2(x+1)][x(x+1) + 3(x+1)][x(x+2) + 3(x+2)] \\ & = (x+1)(x+2)(x+1)(x+3)(x+2)(x+3) \\ & = (x+1)^2(x+2)^2(x+3)^2 \end{aligned}$$

Hence

$$\begin{aligned} & \sqrt{(x^2 + 3x + 2)(x^2 + 4x + 3)(x^2 + 5x + 6)} \\ & = \sqrt{(x+1)^2(x+2)^2(x+3)^2} \\ & = \pm (x+1)(x+2)(x+3) \end{aligned}$$

ix)
$$\begin{aligned} & (x^2 + 8x + 7)(2x^2 - x - 3)(2x^2 + 11x - 21) \\ & = (x^2 + x + 7x + 7)(2x^2 + 2x - 3x - 3)(2x^2 + 14x - 3x - 21) \\ & = [x(x+1) + 7(x+1)][2x(x+1) - 3(x+1)] \\ & \quad [2x(x+7) - 3(x+7)] \\ & = (x+1)(x+7)(x+1)(2x-3)(x+7)(2x-3) \\ & = (x+1)^2(x+7)^2(2x-3)^2 \end{aligned}$$

Hence

$$\begin{aligned} & \sqrt{(x^2 + 8x + 7)(2x^2 - x - 3)(2x^2 + 11x - 21)} \\ & = \sqrt{(x+1)^2(x+7)^2(2x-3)^2} \\ & = \pm (x+1)(x+7)(2x-3) \end{aligned}$$

Q2. Use division method to find the square root of the following expressions.

i)
$$\begin{array}{r} 4x^2 + 12xy + 9y^2 + 16x + 24y + 16 \\ 2x + 3y + 4 \\ \hline 4x^2 + 12xy + 9y^2 + 16x + 24y + 16 \\ 4x^2 \\ \hline 12xy + 9y^2 + 16x + 24y + 16 \\ 12xy + 9y^2 \\ \hline 16x + 24y + 16 \\ 16x + 24y + 16 \\ \hline 0 \end{array}$$

Hence the square root of given expression is
 $\pm (2x + 3y + 4)$

ii)
$$\begin{array}{r} x^4 - 10x^3 + 37x^2 - 60x + 36 \\ x^2 - 5x + 6 \\ \hline x^2 \\ x^4 - 10x^3 + 37x^2 - 60x - 36 \\ -x^4 \\ \hline -10x^3 + 37x^2 - 60x - 36 \\ -10x^3 + 25x^2 \\ \hline 12x^2 - 60x - 36 \\ -12x^2 + 60x - 36 \\ \hline 0 \end{array}$$

Hence $\sqrt{x^4 - 10x^3 + 37x^2 - 60x + 36} = \pm (x^2 - 5x + 6)$

iii) $9x^4 - 6x^3 + 7x^2 - 2x + 1$

$$\begin{array}{r}
 3x^2 - x + 1 \\
 \hline
 3x^2 & 9x^4 - 6x^3 + 7x^2 - 2x + 1 \\
 & -9x^4 \\
 \hline
 6x^2 - x & -6x^4 + 7x^2 - 2x + 1 \\
 & \cancel{-6x^4} \pm x^2 \\
 \hline
 6x^2 - 2x + 1 & 6x^4 - 2x + 1 \\
 & -6x^4 \mp 2x \pm 1 \\
 \hline
 & 0
 \end{array}$$

Hence $\sqrt{9x^4 - 6x^3 + 7x^2 - 2x + 1}$
 $= \pm(3x^2 - x + 1)$

iv) $4 + 25x^2 - 12x - 24x^3 + 16x^4$

In descending order

$$\begin{array}{r}
 = 16x^4 - 24x^3 + 25x^2 - 12x + 4 \\
 4x^2 - 3x + 2
 \end{array}$$

$$\begin{array}{r}
 4x^2 & 16x^4 - 24x^3 + 25x^2 - 12x + 4 \\
 & -16x^4 \\
 \hline
 8x^2 - 3x & -24x^3 + 25x^2 - 12x + 4 \\
 & \cancel{-24x^3} \pm 9x^2 \\
 \hline
 8x^2 - 6x + 2 & 16x^2 - 12x + 4 \\
 & \cancel{-16x^2} \mp 12x \pm 4 \\
 \hline
 & 0
 \end{array}$$

Hence $\sqrt{16x^4 - 24x^3 + 25x^2 - 12x + 4}$
 $= \pm(4x^2 - 3x + 2)$

v) $\frac{x^2}{y^2} - 10\frac{x}{y} + 27 - 10\frac{y}{x} + \frac{y^2}{x^2}$
 $(x \neq 0, y \neq 0)$

Hence

$$\begin{array}{r}
 \frac{x}{y} - 5 + \frac{y}{x} \\
 \hline
 \frac{x}{y} & \frac{x^2}{y^2} - 10\frac{x}{y} + 27 - 10\frac{y}{x} + \frac{y^2}{x^2} \\
 & \cancel{\frac{x^2}{y^2}} \\
 \hline
 \frac{2x}{y} - 5 & -10\frac{x}{y} + 27 - 10\frac{y}{x} + \frac{y^2}{x^2} \\
 & \cancel{-10\frac{x}{y}} \pm 25 \\
 \hline
 \frac{2x}{y} - 10 + \frac{y}{x} & 2 - 10\frac{y}{x} + \frac{y^2}{x^2} \\
 & \cancel{2} \mp 10\frac{y}{x} \pm \frac{y^2}{x^2} \\
 \hline
 & 0
 \end{array}$$

$$\sqrt{\frac{x^2}{y^2} - 10\frac{x}{y} + 27 - 10\frac{y}{x} + \frac{y^2}{x^2}}$$

The required square root

$$= \pm\left(\frac{x}{y} - 5 + \frac{y}{x}\right)$$

Q3. Find the value of 'k' for which the following expression will become a perfect square?

i) $4x^4 - 12x^3 + 37x^2 - 42x + k$

$$\begin{array}{r}
 2x^2 & 2x^2 - 3x + 7 \\
 & 4x^4 - 12x^3 + 37x^2 - 42x + k \\
 & -4x^4 \\
 \hline
 4x^2 - 3x & -12x^3 + 37x^2 - 42x + k \\
 & \cancel{-12x^3} \pm 9x^2 \\
 \hline
 4x^2 - 6x + 7 & 28x^2 - 42x + k \\
 & \cancel{-28x^2} \mp 42x \pm 49 \\
 \hline
 & k - 49
 \end{array}$$

As given that the given expression is a perfect square, so

Remainder = 0

$k - 49 = 0$

$$[k = 49]$$

ii) $x^4 - 4x^3 + 10x^2 - kx + 9$

$$\begin{array}{r} x^2 - 2x + 3 \\ \hline x^2 \left| \begin{array}{r} x^4 - 4x^3 + 10x^2 - kx + 9 \\ -x^4 \\ \hline -4x^3 + 10x^2 - kx + 9 \\ -4x^3 \pm 4x^2 \\ \hline 6x^2 - kx + 9 \\ 6x^2 \mp 12x \pm 9 \\ \hline (-k+12)x \end{array} \right. \end{array}$$

As given that the given expression is a perfect square, so

$$\text{Remainder} = 0$$

$$(-k+12)x = 0$$

As $x \neq 0$, so $-k+12=0$

$$\Rightarrow [k = 12]$$

Q4. Find the values of ' l ' and ' m ' for which the following expression will become perfect square.

i) $x^4 + 4x^3 + 16x^2 + lx + m$

$$\begin{array}{r} x^2 + 2x + 6 \\ \hline x^2 \left| \begin{array}{r} x^4 + 4x^3 + 16x^2 + lx + m \\ -x^4 \\ \hline 4x^3 + 16x^2 + lx + m \\ -4x^3 \pm 4x^2 \\ \hline 12x^2 + lx + m \\ -12x^2 \pm 24x \pm 36 \\ \hline (l-24)x + (m-36) \end{array} \right. \end{array}$$

As the given expression is to be a perfect square, so

$$\text{Remainder} = 0$$

$$(l-24)x + (m-36) = 0$$

As $x \neq 0$, so $l-24=0$ and $m-36=0$

$$\Rightarrow [l = 24] \text{ and } [m = 36]$$

ii) $49x^4 - 70x^3 + 109x^2 + lx - m$

$$\begin{array}{r} 7x^2 \left| \begin{array}{r} 7x^2 - 5x + 6 \\ \hline 49x^4 - 70x^3 + 109x^2 + lx - m \\ -49x^4 \\ \hline -70x^3 + 109x^2 + lx - m \\ -70x^3 \pm 25x^2 \\ \hline 84x^2 + lx - m \\ -84x^2 \mp 60x \pm 36 \\ \hline (l+60)x - m - 36 \end{array} \right. \end{array}$$

As the given expression is to be a perfect square, so

$$(l+60)x - m - 36 = 0$$

As $x \neq 0$, so $l+60=0$ and $-m-36=0$

$$\Rightarrow [l = -60] \text{ and } [m = -36]$$

Q5. To make the expression

$9x^4 - 12x^3 + 22x^2 - 13x + 12$ a perfect square.

i) What should be added to it?

ii) What should be subtracted from it?

iii) What should be the value of ' x '?

$$\begin{array}{r} 3x^2 \left| \begin{array}{r} 3x^2 - 2x + 3 \\ \hline 3x^2 \left| \begin{array}{r} 9x^4 - 12x^3 + 22x^2 - 13x + 12 \\ -9x^4 \\ \hline -12x^3 + 22x^2 - 13x + 12 \\ -12x^3 \pm 4x^2 \\ \hline 18x^2 - 13x + 12 \\ -18x^2 \mp 12x \pm 9 \\ \hline -x + 3 \end{array} \right. \end{array} \right. \end{array}$$

To make the given expression a complete square

i) $x-3$ should be added

ii) $-x+3$ should be subtracted

iii) For value of 'x'

$$\begin{aligned}\text{Remainder} &= 0 \\ -x + 3 &= 0 \\ \boxed{x = 3}\end{aligned}$$

Q6. Find H.C.F of following by factorization

$$8x^4 - 128, 12x^3 - 96.$$

Solution:

$$\begin{aligned}8x^4 - 128 &= 8(x^4 - 16) \\ &= 8((x^2)^2 - (4)^2) \\ &= 8(x^2 + 4)(x^2 - 4) \\ &= 8(x^2 + 4)(x+2)(x-2) \\ 12x^3 - 96 &= 12(x^3 - 8) \\ &= 12(x^3 - 2^3) \\ &= 12(x-2)(x^2 + 2x + 4)\end{aligned}$$

$$\begin{array}{ll}\text{Common factor} & = 4(x-2) \\ \text{H.C.F} & = 4(x-2)\end{array}$$

Q7. Find H.C.F of following by division method.

$$y^3 + 3y^2 - 3y - 9, y^3 + 3y^2 - 8y - 24$$

Solution:

1

$$\begin{array}{r}y^3 + 3y^2 - 3y - 9 \quad y^3 + 3y^2 - 8y - 24 \\ \underline{-y^3 - 3y^2 - 3y - 9} \\ -5y - 15 \\ -5(y+3) \\ \quad\quad\quad y^2 - 3 \\ (y+3) \quad y^3 + 3y^2 - 3y - 9 \\ - y^3 - 3y^2 \\ - 3y - 9 \\ + 3y + 9 \\ \quad\quad\quad x\end{array}$$

$$\text{H.C.F} = y + 3$$

Q8. Find L.C.M of following by factorization.

$$12x^2 - 75, 6x^2 - 13x - 5, 4x^2 - 20x + 25$$

Solution:

$$\begin{aligned}12x^2 - 75 &= 3(4x^2 - 25) \\ &= 3((2x)^2 - (5)^2) \\ &= 3(2x+5)(2x-5) \\ 6x^2 - 13x - 5 &= 6x^2 - 15x + 2x - 5 \\ &= 3x(2x-5) + 1(2x-5)\end{aligned}$$

$$\begin{aligned}4x^2 - 20x + 25 &= (2x)^2 + (5)^2 - 2(2x)(5) \\ &= (2x-5)^2 \\ &= (2x-5)(2x-5)\end{aligned}$$

$$\begin{aligned}\text{L.C.M} &= (2x-5)^2 \times 3(2x+5)(3x+1) \\ &= 3(2x-5)^2(2x+5)(3x+1)\end{aligned}$$

Q9. If H.C.F of $x^4 + 3x^3 + 5x^2 + 26x + 56$ and $x^4 + 2x^3 - 4x^2 - x + 28$ is $x^2 + 5x + 7$, find the

Solution:

$$\text{L.C.M} = \frac{(x^4 + 3x^3 + 5x^2 + 26x + 56)(x^4 + 2x^3 - 4x^2 - x + 28)}{x^2 + 5x + 7}$$

$$\begin{array}{r}x^4 + 3x^3 + 5x^2 + 26x + 56 \\ \underline{-x^4 - 5x^3 - 7x^2} \\ \hline -2x^3 - 2x^2 + 26x + 56 \\ \underline{-2x^3 - 10x^2 - 14x} \\ \hline 8x^2 + 40x + 56\end{array}$$

$$\begin{array}{r}8x^2 + 40x + 56 \\ \underline{-8x^2 - 40x - 56} \\ \hline \times\end{array}$$

L.C.M

$$= (x^2 - 2x + 8)(x^4 + 2x^3 - 4x^2 - x + 28)$$

Q10. Simplify

$$\begin{aligned}\text{(i)} \quad & \frac{3}{x^3 + x^2 + x + 1} - \frac{3}{x^3 - x^2 + x - 1} \\ & \frac{3}{(x^2 + 1)(x + 1)} - \frac{3}{(x^2 + 1)(x - 1)} \\ & = \frac{3(x-1) - 3(x+1)}{(x^2 + 1)(x+1)(x-1)} \\ & = \frac{3x - 3 - 3x - 3}{(x^2 + 1)(x+1)(x-1)} \\ & = \frac{-6}{(x^2 + 1)(x+1)(x-1)} \\ & = \frac{-6}{(x^2 + 1)(x^2 - 1)}\end{aligned}$$

$$= \frac{-6}{x^4 - 1} = \frac{6}{1-x^4} \text{ Ans.}$$

$$\begin{aligned} \text{(ii)} \quad & \frac{a+b}{a^2-b^2} \div \frac{a^2-ab}{a^2-2ab+b^2} \\ &= \frac{a+b}{(a-b)(a+b)} \div \frac{a(a-b)}{(a-b)^2} \\ &= \frac{1}{a-b} \div \frac{a}{a-b} \\ &= \frac{1}{a-b} \times \frac{a-b}{a} \\ &= \frac{1}{a} \end{aligned}$$

Q11. Find square root by using factorization

$$\left(x^2 + \frac{1}{x^2}\right) + 10\left(x + \frac{1}{x}\right) + 27 \quad (x \neq 0)$$

Solution:

$$= \left(x^2 + \frac{1}{x^2}\right) + 10\left(x + \frac{1}{x}\right) + 25 + 2$$

Q12. Find square root by using division method.

$$\frac{4x^2}{y^2} + \frac{20x}{y} + 13 - \frac{30y}{x} + \frac{9y^2}{x^2} \quad (x, y \neq 0)$$

Solution:

$\frac{2x}{y} + 5 - \frac{3y}{x}$ $\frac{2x}{y}$ $\frac{4x^2}{y^2} + \frac{20x}{y} + 13 - \frac{30y}{x} + \frac{9y^2}{x^2}$ $\frac{4x^2}{y^2}$ \hline $\frac{4x}{y} + 5$	$\frac{20}{y}x + 13$ $\frac{20}{y}x \pm 25$ \hline $\frac{4x}{y} + 10 - \frac{3y}{x}$ $-12 - \frac{30y}{x} + \frac{9y^2}{x^2}$ $\mp 12 \mp \frac{30y}{x} \pm \frac{9y^2}{x^2}$ \hline \times
--	---

$$\text{Required square root} = \pm \left(\frac{2x}{y} + 5 - \frac{3y}{x} \right)$$

$$\begin{aligned} &= x^2 + \frac{1}{x^2} + 2 + 10\left(x + \frac{1}{x}\right) + 25 \\ &= \left(x + \frac{1}{x}\right)^2 + 10\left(x + \frac{1}{x}\right) + 25 \\ &\text{Let } x + \frac{1}{x} = a \\ &= a^2 + 10a + 25 \\ &= (a+5)^2 \\ &\text{Taking square root} \\ &= \sqrt{(\pm(a+5))^2} \\ &= \pm(a+5) \\ &= \pm\left(x + \frac{1}{x} + 5\right) \end{aligned}$$

Objective

1. H.C.F of $p^3q - pq^3$ and $p^5q^2 - p^2q^5$ is _____
 - $pq(p^2 - q^2)$
 - $pq(p - q)$
 - $p^2q^2(p - q)$
 - $pq(p^3 - q^3)$
2. H.C.F. of $5x^2y^2$ and $20x^3y^3$ is: _____
 - $5x^2y^2$
 - $20x^3y^3$
 - $100x^5y^5$
 - $5xy$
3. H.C.F of $x - 2$ and $x^2 + x - 6$ is _____
 - $x^2 + x - 6$
 - $x + 2$
 - $x - 2$
 - $x + 2$
4. H.C.F of $a^3 + b^3$ and $a^2 - ab + b^2$ is _____
 - $a + b$
 - $a^2 - ab + b^2$
 - $(a-b)^2$
 - $a^2 + b^2$
5. H.C.F of $x^2 - 5x + 6$ and $x^2 - x - 6$ is _____:
 - $x - 3$
 - $x + 2$
 - $x^2 - 4$
 - $x - 2$
6. H.C.F of $a^2 - b^2$ and $a^3 - b^3$ is _____
 - $a - b$
 - $a + b$
 - $a^2 + ab + b^2$
 - $a^2 - ab + b^2$
7. H.C.F of $x^2 + 3x + 2$, $x^2 + 4x + 3$, $x^2 + 5x + 4$ is:
 - $x+1$
 - $(x+1)(x+2)$
 - $(x+3)$
 - $(x+4)(x+1)$
8. L.C.M of $15x^2$, $45xy$ and $30xyz$ is _____
 - $90xyz$
 - $90x^2yz$
 - $15xyz$
 - $15x^2yz$
9. L.C.M of a^2+b^2 and a^4-b^4 is:
 - $a^2 + b^2$
 - $a^2 - b^2$
 - $a^4 - b^4$
 - $a - b$
10. The product of two algebraic expression is equal to the _____ of

- their H.C.F and L.C.M.
- Sum
 - Difference
 - Product
 - Quotient
11. Simplify $\frac{a}{9a^2 - b^2} + \frac{1}{3a - b} = \underline{\hspace{2cm}}$
 - $\frac{4a}{9a^2 - b^2}$
 - $\frac{4a - b}{9a^2 - b^2}$
 - $\frac{4a + b}{9a^2 - b^2}$
 - $\frac{b}{9a^2 - b^2}$
 12. Simplify $\frac{a^2 + 5a - 14}{a^2 - 3a - 18} \times \frac{a+3}{a-2} = \underline{\hspace{2cm}}$
 - $\frac{a+7}{a-6}$
 - $\frac{a+7}{a-2}$
 - $\frac{a+3}{a-6}$
 - $\frac{a-3}{a+2}$
 13. Simplify

$$\frac{a^3 - b^3}{a^4 - b^4} \div \left(\frac{a^2 + ab + b^2}{a^2 + b^2} \right) = \underline{\hspace{2cm}}$$
 - $\frac{1}{a+b}$
 - $\frac{1}{a-b}$
 - $\frac{a-b}{a^2+b^2}$
 - $\frac{a+b}{a^2+b^2}$
 14. Simplify :

$$\left(\frac{2x+y}{x+y} - 1 \right) \div \left(1 - \frac{x}{x+y} \right) = \underline{\hspace{2cm}}$$

- (a) $\frac{x}{x+y}$ (b) $\frac{x}{x-y}$
 (c) $\frac{y}{x}$ (d) $\frac{x}{y}$

15. The square root of $a^2 - 2a + 1$ is ____
 (a) $\pm(a+1)$ (b) $\pm(a-1)$
 (c) $a-1$ (d) $a+1$
16. What should be added to complete the square of $x^4 + 64$?
 (a) $8x^2$ (b) $-8x^2$
 (c) $16x^2$ (d) $4x^2$
17. The square root of $x^4 + \frac{1}{x^4} + 2$ is
 —
 (a) $\pm\left(x + \frac{1}{x}\right)$ (b) $\pm\left(x^2 + \frac{1}{x^2}\right)$
 (c) $\pm\left(x - \frac{1}{x}\right)$ (d) $\pm\left(x^2 - \frac{1}{x^2}\right)$
18. The square root of $4x^2 - 12x + 9$ is:
 (a) $\pm(2x - 3)$
 (b) $\pm(2x + 3)$
 (c) $(2x + 3)^2$
 (d) $(2x - 3)^2$

19. L.C.M = ____
 (a) $\frac{p(x) \times q(x)}{\text{H.C.F}}$ (b) $\frac{p(x) \cdot q(x)}{\text{L.C.M}}$
 (c) $\frac{p(x)}{q(x) \times \text{H.C.F}}$ (d) $\frac{q(x)}{p(x) \times \text{H.C.F}}$
20. H.C.F. = ____
 (a) $\frac{p(x) \times q(x)}{\text{L.C.M}}$ (b) $\frac{p(x) \times q(x)}{\text{H.C.F}}$
 (c) $\frac{p(x)}{q(x) \times \text{L.C.M}}$ (d) $\frac{\text{L.C.M}}{p(x) \times q(x)}$
21. L.C.M x H.C.F =
 (a) $p(x) \times q(x)$ (b) $p(x) \times \text{H.C.F}$
 (c) $q(x) \times \text{L.C.M}$ (d) None
22. Any unknown expression may be found if ____ of them are known by using the relation
 $\text{L.C.M} \times \text{H.C.F} = p(x) \times q(x)$
 (a) Two
 (b) Three
 (c) Four
 (d) None

ANSWER KEY

1.	a	2.	a	3.	c	4.	b	5.	a
6.	a	7.	a	8.	b	9.	c	10.	c
11.	c	12.	a	13.	a	14.	d	15.	b
16.	c	17.	b	18.	a	19.	a	20.	a
21.	a	22.	b						